

ONMedU, Department of Obstetrics and Gynecology. Practical class №13. Physiology of labor.
Analgesia of labor.

**MINISTRY OF HEALTH OF UKRAINE
ODESA NATIONAL MEDICAL UNIVERSITY**

International Faculty

Department of obstetrics and gynecology



CONFIRMED by
Vice-rector for scientific and
pedagogical work
Eduard BURIACHKIVSKYI
September 1, 2023

**METHODOLOGICAL RECOMMENDATIONS
FOR PRACTICAL CLASS**

International Faculty, Course V

Discipline "Obstetrics and Gynecology"

Practical lesson №13. Topic: Physiology of labor. Analgesia of labor.

Methodological recommendations for practical lesson. «Health care», master's degree in the
specialty "Medicine". Discipline "Obstetrics and Gynecology"

ONMedU, Department of Obstetrics and Gynecology. Practical class №13. Physiology of labor.
Analgesia of labor.

Approved:

Meeting of the Department of Obstetrics and Gynecology of Odesa National
Medical University


Protocol No. 1 dated August 28, 2023

Head of the Department  (Ihor GLADCHUK)

Developer:

Ph.D., assistant professor of

obstetrics and gynecology department

 (O.Nadvorna)

Methodological recommendations for practical lesson. «Health care», master's degree in the
specialty "Medicine". Discipline "Obstetrics and Gynecology"

Practical class №13.

PHYSIOLOGY OF LABOUR. ANALGESIA OF LABOUR.

LEARNING OBJECTIVE. Labour is a sequence of uterine contractions that results in effacement and dilatation of the cervix and voluntary bearing-down efforts leading to the expulsion per vagina of the products of conception. Delivery is the mode of expulsion of the fetus and placenta. Labour and delivery is a normal physiologic process that most women experience without complications. The goal of the management of this process is to foster a safe birth for mothers and their newborns. Additionally, the staff should attempt to make the patient and her support person(s) feel welcome, comfortable, and informed throughout the labour and delivery process. Physical contact between the newborn and the parents in the delivery room should be encouraged. Every effort should be made to foster family interaction and to support the desire of the family to be together. The role of the obstetrician/midwife and the labour and delivery staff is to anticipate and manage complications that may occur that could harm the mother or the fetus. When a decision is made to intervene, it must be considered carefully, because each intervention carries both potential benefits and potential risks. The best management in the majority of cases may be close observation and, when necessary, cautious intervention.

Obstetric analgesia or anesthesia refers to the multiple techniques useful for the alleviation of pain associated with labour, delivery, or surgery. The choice of an appropriate analgesic technique must be made by the patient, the obstetrician, and the anesthesiologist and should take into consideration the patient's anatomy and physiology, the status of her fetus, the obstetric plan for delivery, and the pharmacology of the drugs to be employed.

BASIC CONCEPTS: Labour is a sequence of uterine contractions that results in effacement and dilatation of the cervix and voluntary bearing-down efforts leading to the expulsion per vagina of the products of conception.

EQUIPMENT

- Multimedia equipment (computer, projector, screen), TV.
- Obstetric models and obstetric instruments (pelvimeter, obstetric stethoscope, centimeter tape).
- Professional algorithms, structural-logical schemes, tables, videos.
- Results of laboratory and instrumental researches, situational tasks, patients, medical histories.

EDUCATIONAL TIME – 4 h

1. ORGANIZATIONAL STAGE

- Greetings,

- checking attendees,
- defining of educational goals,
- providing of positive motivation.

Delivery is the mode of expulsion of the fetus and placenta. Labour and delivery is a normal physiologic process that most women experience without complications. The goal of the management of this process is to foster a safe birth for mothers and their newborns. Additionally, the staff should attempt to make the patient and her support person(s) feel welcome, comfortable, and informed throughout the labour and delivery process. Physical contact between the newborn and the parents in the delivery room should be encouraged.

2. CONTROL OF BASIC KNOWLEDGE (written work, written testing, online testing, face-to-face interview, etc.)

2.1. Requirements for the theoretical readiness of students to perform practical classes.

Knowledge requirements:

- Communication and clinical examination skills.
- Ability to determine the list of required clinical, laboratory and instrumental studies and evaluate their results.
- Ability to make a preliminary and clinical diagnosis of the labour
- Ability to perform medical manipulations
- Ability to determine the tactics of physiological pregnancy, physiological labour and the postpartum period.
- Ability to keep medical records.

List of didactic units:

- Pelvis from anatomical and obstetric points of view.
- Pelvic floor.
- The structure of the fetal head.
- The dimensions of the fetal head and body.
- Signs of fetal maturity.
- Measurement and evaluation of the pelvis
- Determination of the stage of labour, period and phase of labour.
- Signs of placentas separation in the third period of labour

2.2. Questions (test tasks, tasks, clinical situations) to test basic knowledge on the topic of the class.

Questions:

- Evaluation of the cervical status (scale Bishop),
- How to determine the beginning of the first period of delivery, objectively assess the character of labour activity (degree of cervical effacement, frequency, strength and duration of the contractions),
- How to fill in f partogram,
- How to identify and assess fetal heart (auscultation, cardiotocogram),
- Obstetric analgesia and anesthesia,
- Active management of the third stage of labour (demonstration on the phantom),
- Expectant management of the third stage of labour (demonstration on the phantom),
- Examination of the placenta,
- Determination the total blood loss during delivery,
- Essentials of normal newborn assessment (Apgar scores).

Test tasks

Direction: For each of the multiple-choice questions select the lettered answer that is the one best response in each case.

1. One of the planes of the pelvis is limited behind sacral promontory, in front – crista iliaca and superior edge of the pubis symphysis, on each side - lin. terminalis.

What moment of the biomechanism of birth does the fetus make in the given pelvic plane?

- A. It does not make any of the moments.
- B. External turn of the head and internal turn of the shoulders.
- C. Extension of the head.
- D. Internal turn of the head.
- E. Flexion of the head.

2. On the occipital area of the head of a newborn, having a dolichocephalic form, there is a patrimonial tumour with the centre at the area of the small fontanel. What head presentation did the birth occur in?

- A. Frontal.
- B. Anterior type, occipital presentation.
- C. Facial.
- D. Posterior type, occipital presentation.
- E. Frontoparietal.

3. During vaginal examination 6 hours after the beginning of patrimonial activity the following was revealed: the cervix is open 5 cm, the fetal head is presented and is pressed to the input of the pelvis. The sagittal suture is in the right diagonal diameter,

the small fontanel is to the left, to the side. What moment of the biomechanism of birth is being described?

- A. Flexion of the head.
- B. Extension of the head.
- C. Internal turn of the head.
- D. Additional flexion of the head.
- E. Internal turn the shoulders.

4. During vaginal examination, the fetal head is determined; it fills the posterior surface of the symphysis pubis and hollow of the sacrum, a inferior edge of the symphysis pubis, spines of the ischiadic bones, sacrococcygeal joints are accessible with palpation. In what plane of the small pelvis is the head located?

- A. Plane of the narrow part of the pelvic cavity.
- B. Plane of the wide part of the pelvic cavity.
- C. On the input to the pelvis.
- D. In the input to the pelvis.
- E. In the exit of the pelvis.

5. Secundipara has been in childbirth for 8 hours. Light amniotic fluid was discharged. The fetal position is longitudinal, the fetal head is not determined over the input to the pelvis. Fetal palpitation is clear, rhythmical, 140 b.p.m., over the pubis. Internal obstetrical examination: the cervix is smooth, fully open, the fetal sac is absent. The hollow of the sacrum is completely filled with the head. Spina ischiadica is not reached. The sagittal suture is in the direct diameter of the pelvis. The large fontanel is at the pubis. Contractions have begun. What period of birth is being described?

- A I period.
- B. End of the I period.
- C. Beginning of the II period.
- D. End of the II period.
- E. Beginning of the III period.

6. A woman in her first pregnancy had a live baby boy, weight 3200 g, length 50 cm. The umbilical cord was transected after pulsation of the vessels stopped. When the edge of the palm is pressed on the symphysis, the umbilical cord retreats into the vagina. What sign is used for determining whether the placenta has separated from the uterus?

- A. Alfelda.
- B. Kustner-Chukalova.
- C. Schröder.
- D. Dovjenko.
- E. Rogovina.

7. Primipara gave birth to a live mature baby boy with an estimation by the Apgar scale of 9 points. At the present moment, the umbilical cord vessels are not pulsating, the umbilical cord was cut. Bloody discharge from the genital tract are absent. Specify the period of birth:

- A. Early postnatal period.
- B. Opening.
- C. Expulsion of fetus.
- D. Placental stage.
- E. Late postnatal period.

8. Data from the internal obstetrical examination of a parturient woman: the cervix uterus is smooth, open 6 cm. The fetal sac is intact. The fetal head is presented, when pressed on, it does not make push back. During palpation of the pelvic, the pubis symphysis, innominate lines, promontory are free. The sagittal suture is in the transverse diameter, the small fontanel is to the right at the pubis. In what plane is the fetal head?

- A. Pressed to the input of the pelvis.
- B. On the input of the pelvis.
- C. Wide part of the pelvic cavity.
- D. Narrow part of the pelvic cavity.
- E. Exit of the pelvis.

9. I period of I due labour. Fetal position - longitudinal, the small segment of the head is in the input to the pelvis. Fetal palpitation is clear, rhythmical, 140 b.p.m., to the left below the navel. During internal obstetrical examination, the cervix is smooth, open 6 cm, the small segment of the fetal head is in the input to the pelvis. The sagittal suture is in the right diagonal diameter, the small fontanel is to the left closer to the pubis. Specify the position and type of position:

- A. High direct standing of the sagittal suture.
- B. I position, posterior type of position.
- C. II position, anterior type of position.
- D. II position, posterior type of position.
- E. I position, anterior type of position.

10. Secundipara was delivered to the maternity hospital 6 hours after the beginning of labour. Contractions are 30-35 seconds, every 4 minutes, good force. BP is 120/80 mm hg. Pulse is 80 b.p.m., rhythmical, of satisfactory properties. The fetal heart beat is 146 b.p.m. The fetal position is longitudinal, head presentation, I

position, anterior type of position. The small segment of the fetal head is in the input to the pelvis. The height of the contraction ring is 5 cm over the pubis. When should internal obstetrical examination be performed?

- A. Upon arrival and after amniotic discharge.
- B. Every 2 hours.
- C. At the beginning of the II period of birth.
- D. When transferred to the postnatal ward.
- E. At the end of the I period of birth.

11. In what size of the plane of the input to the pelvis is the sagittal suture during anterior position, occipital presentation, I position after performing flexion of the head?

- A. Direct.
- B. Left diagonal or direct.
- C. Right diagonal or direct.
- D. Right diagonal or transverse.
- E. Left diagonal or transverse.

12. What size of the plane of the input to the pelvis is the sagittal suture in during posterior type of position, occipital presentation, I position after flexion of the head?

- A. Direct.
- B. Left diagonal or direct.
- C. Right diagonal or direct.
- D. Right diagonal or transverse.
- E. Left diagonal or transverse.

13. In what plane of the pelvis does the fetus finishing making the internal turn of its head during anterior type of position, occipital presentation, II position?

- A. Input to the pelvis.
- B. The wide part of the pelvic cavity.
- C. The narrow part of the pelvic cavity.
- D. When going from the wide part of the cavity to the narrow part of the pelvic cavity.
- E. The exit of the pelvis.

14. In what diameter of the plane of the exit from the pelvis is the sagittal suture during extension of the head during posterior type of position, occipital presentation, I position?

- A. Direct.
- B. Left diagonal or direct.
- C. Right diagonal or direct.
- D. Right diagonal or transverse.
- E. Left diagonal or transverse.

15. In what diameter of the plane of the exit from the pelvis does the fetal shoulders cut through and are born during the anterior type of position, occipital presentation?

- A. Direct.
- B. Right diagonal.
- C. Left diagonal.
- D. Transverse.
- E. Direct and transverse.

16. The head of a newborn has a dolichocephalic form, extended from the front to the back. At the occipital area, there is a patrimonial tumour located in the middle between the large and small fontanel. What head presentation did the birth occur in?

- A. Frontoparietal.
- B. Anterior type, occipital presentation.
- C. Posterior type, occipital presentation.
- D. Facial.
- E. Frontal.

17. Primipara, 20 years old, is in the beginning of the I period of physiological birth. Contractions are 15-20 sec., every 10-15 min., weak. Fetal palpitation is not suffering. How much in cm should the cervix be open for the amniotic discharge to be timely?

- A. 6-8.
- B. 8-10.
- C. 4-6.
- D. 2-4.
- E. 1,5-2.

18. During internal obstetrical examination of a secundipara, the cervix is open 4-5 cm, when pressing with a finger on the fetal head, it pushes away and again returns to its initial position, the pelvis is free. How long is the fetal head permitted to be in this plane, in hours?

- A. 5.
- B. 6.
- C. 4.
- D. 3.
- E. 2.

19. You are observing a parturient woman in the 1 period of birth. During internal obstetrical examination, the sagittal suture is in the right diagonal diameter, the small fontanel is closer to the pubis. How should the parturient woman lay in bed, so that the fetal head would be insert correctly?

- A. On her left side.
- B. On her back.
- C. On her right side.
- D. It make do difference.
- E. She is only allowed to walk.

20. Secundipara with a weight of 80 kg. What is the acceptable amount of blood loss, ml:

- A. 400.
- B. 500.
- C. 600.
- D. 700.
- E. 800.

Answer key

1.	E	12.	E	23.	
2.	D	13.	E	24.	
3.	A	14.	A	25.	
4.	A	15.	A	26.	
5.	C	16.	B	27.	
6.	B	17.	B	28.	
7.	D	18.	B	29.	
8.	A	19.	A	30.	
9.	E	20.	A	31.	
10.	A	21.		32.	
11.	D	22.		33.	

3. FORMATION OF PROFESSIONAL SKILLS (mastering skills, conducting curation, determining the treatment regimen, conducting a laboratory study, etc.).

3.1. Content of tasks (tasks, clinical situations, etc.).

Interactive task:

Students of the group are divided into 3 subgroups of 3-4 people each. They work in the classroom, reception department of the maternity hospital, labor & delivery ward, neonatal department with pregnant and newborns.

Tasks:

Subgroup I - to perform maneuvers of external obstetrical examination, (Leopold's maneuvers). Define presentation, position, type of fetus. Demonstrate on the phantom active tactics III stage of labour. Demonstrate by the phantom expectant tactics III stage of labour.

Subgroup II - Compare and contrast the advantages and disadvantages of external and internal fetal monitoring, including the appropriate use for each. Estimated date of birth, to determine if term or preterm. Fetal movement (frequency in the past few days)

- to assess progress and stage of labour provided by partogram.
- the status of the newborn Apgar scores (in case of demonstrations delivery) grade

Subgroup III – to assess answers of subgroups I and II and makes adjustments.

Tests:

Direction: For each of the multiple- choice questions select the lettered answer that is the one best response in each case.

1. In what size of the plane of the input to the pelvis is the sagittal suture during anterior position, occipital presentation, I position after performing flexion of the head?

- A. Direct.
- B. Left diagonal or direct.
- C. Right diagonal or direct.
- D. Right diagonal or transverse.
- E. Left diagonal or transverse.

2. What size of the plane of the input to the pelvis is the sagittal suture in during posterior type of position, occipital presentation, I position after flexion of the head?

- A. Direct.
- B. Left diagonal or direct.
- C. Right diagonal or direct.
- D. Right diagonal or transverse.
- E. Left diagonal or transverse.

3. In what plane of the pelvis does the fetus finishing making the internal turn of its head during anterior type of position, occipital presentation, II position?

- A. Input to the pelvis.
- B. The wide part of the pelvic cavity.
- C. The narrow part of the pelvic cavity.
- D. When going from the wide part of the cavity to the narrow part of the pelvic cavity.
- E. The exit of the pelvis.

4. In what diameter of the plane of the exit from the pelvis is the sagittal suture during extension of the head during posterior type of position, occipital presentation, I position?

- A. Direct.
- B. Left diagonal or direct.
- C. Right diagonal or direct.
- D. Right diagonal or transverse.
- E. Left diagonal or transverse.

5. In what diameter of the plane of the exit from the pelvis does the fetal shoulders cut through and are born during the anterior type of position, occipital presentation?

- A. Direct.
- B. Right diagonal.
- C. Left diagonal.
- D. Transverse.
- E. Direct and transverse.

Answer key

1.	D
2.	E
3.	E
4.	A
5.	A

3.2. Educational materials, recommendations (instructions) for performing tasks

The course of physiology normal labour and delivery comprises a complex relationship between several dynamic parameters, including uterine contractions, cervical dilation, fetal descent, and elapsed time. The onset of labour in humans occurs around 280 days, or 37-42 weeks, from the first day of a patient's last menstrual period (LMP).

1. Precursors of birth:

- 1.1.the lower the bottom of the uterus,
- 1.2.increase response of the uterine to mechanical stimuli,
- 1.3.going out mucus plug of the cervical canal,
- 1.4.loss of women weight 1-1,5 kg,
- 1.5.reduce the amount of amniotic fluid,
- 1.6.fetal position; the orientation of the presenting vertex within the maternal pelvis.

2. Physiological preliminary period lasts about 6 hours and is characterized by irregular increasing in frequency, duration and intensity bouts of pain of cramping nature, mostly in the abdomen and in the groin. Pregnant women do not feel tired, their sleep is not disturbed. Irregular uterine contractions get intensified and gradually turn into regular maternity activity (contractions). In the intervals between contractions the uterine tone is not intensified, the fetal heart beat is clear and smooth, the cervix is mature, vaginal discharge is "mucous plug" mixed with blood.

3. Readiness for labour.

Labour is the physiological process by which the fetus and the placenta are expelled from the uterus through the birth canal.

Beginning of labour is an appearing of 2 contractions during 10 min. Duration of each contraction 20 seconds or more. Labour is not an even process. Within the first 4 – 5 h complex structural changes in cervix occur: it becomes shorter, smoothes, merges with lower segment of the uterus. The general length of I stage of labour during the management of labour act is 10 – 12 h in nulliparous women and 6 – 8 h in multiparous women.

4. Identifying readiness of the organism of a woman for childbirth.

Readiness of the female organism for childbirth is determined by examining the characteristic changes occurring in the cervix.

- 4.1. cervical conditon,
- 4.2. length of cervix (cm),
- 4.3. cervical dilatation (cm),
- 4.4. position of cervix,
- 4.5. station of presenting part.

There are 3 grades of cervical state: immature, not fully mature and mature cervix. Bishop Scoring System

Factor		Points		
		0	1	2
1.	Position of cervix	Directed toward the symphysis	Middle	The pelvic axis
2.	Length of cervix (cm)	> 2 cm	1-2 cm	< 1 cm
3.	Consistency of cervix	Dense	Mederate	Soft
4.	Cervical dilatation (cm)	close	1-2	> 2

5.	Station of presenting part	above the pelvic inlet plane	Between the superior and posterior margin of the symphysis	posterior margin of the symphysis and below.
-----------	-----------------------------------	------------------------------	--	--

0-2 points – «immature» cervix;

3-5 points – «not fully mature» cervix;

> 6 points – «mature» cervix.

5. Cardinal Movements of Labour

From the perspective of the fetus, labour involves movement progressively downward through the pelvis by the following cardinal movements.

5. 1. Mechanisms of labour with occiput anterior presentation. There are four moments.

First moment – flexion of the fetus head,

Second moment – descent and internal rotation,

Third moment – extension of the fetus head,

Fourth moment – external rotation.

5.2. Mechanisms of labour with occiput posterior presentation. It consists of five moments.

First moment – flexion of the fetus head,

Second moment – descent and internal rotation,


Third moment – additional flexion of the fetal head,

Fourth moment – extension of the fetus head,

Fifth moment – external rotation.

6. Regulation of labour activity

In late pregnancy and early childbirth a woman body is dominated by processes of inhibition of the cerebral cortex and increase in the excitability of the subcortical structures of the spinal cord. By this time a pregnant woman has shaped structure of the childbirth dominant, which is characterized by increased reaction to interoceptive stimuli of the cervix uteri and the weakening or complete absence of reaction to extraceptive stimuli, which is the prerequisite for a normal, uncomplicated course of the generic process.



The diagram of regulation of uterine activity. solid arrows - activation (stimulation), dashed – suppression.

7. The clinical course of childbirth

There are 3 stages of labour:

I—dilation of the cervix,

II—expulsion of the fetus,

III— postnatal stage.

Diagnosis periods and phases of delivery:

Symptoms and signs	Period	Phase
Cervix not opened	False labour / Absence of labour activity /	
Cervix opened less than 3 cm in primipara, 4 cm for multiparous	First	Latent
Cervix is opened at 3-9 cm.	The rate of opening of the cervix least (or more) - 1 cm / hour.	
Start descent the fetal head	First	Active
Full of cervical effacement (10 cm).	The head of the fetus is in the pelvic cavity.	
No urge to bearing-down	Second	Early

Methodological recommendations for practical lesson. «Health care», master's degree in the specialty "Medicine". Discipline "Obstetrics and Gynecology"

Full of cervical effacement (10 cm).	The presenting part of fetus reaching the pelvic bottom (plane of the pelvic outlet).	
To start to have bearing-down	Second	Late (bearing-down pains)
The third stage of labour begins after delivery of the newborn and ends with the expulsion of placenta.	Third	

7.1. The first stage of labour consists of the time between the onset of regular contractions associated with cervical change and the occurrence of complete cervical dilation (10 cm).

The first stage is further divided into latent and active phases. Although the distinction between the two phases can be difficult to make, the latent phase of labour is characterized by a slower rate of cervical dilation despite strong, regular uterine contractions.

7.1.1. *The latent phase* can normally last up to 4-5 hours in multigravid patients and up to 6-8 hours in nulligravidas. In the active phase of labour, there is a more rapid change in cervical dilation. Patients may move extremely rapidly through active labour, although the lower limit of normal for cervical change is about 1 cm per hour for nulliparous women.

7.1.2. *Active phases.* In the majority of patients, the transition between the latent and active phases occurs at some time between 3 and 5 cm of cervical dilation, although it is possible, particularly in multigravid patients, to see a patient who is 5-cm dilated and still in the latent phase of labour. It is also critical for the clinician to accurately distinguish between latent phase labour, during which incremental cervical change is occurring (although slowly), and dysfunctional uterine contractions, a condition characterized by no change in cervical dilation despite strong, painful uterine contractions. Such dysfunctional contractions do not constitute labour, and treating them as such may lead to unnecessary intervention.

Characteristics contractions the active phase I period of labour:

- strength - moderate,
- lasting - 45-50 seconds
- frequency - 3-5 contractions in 10 minutes.

7.2. The second stage of labour (expulsion of the fetus).

The second stage of labour is defined as the interval between complete cervical dilation and delivery of the baby. This stage is characterized by descent of the fetal presenting part; maternal sensation of pelvic pressure as this descent progresses; and maternal expulsive efforts, which in concert with uterine contractions effect delivery of the baby. The duration of the second stage varies with parity, ethnicity, fetal size, and the presence or absence of regional anesthesia and can range from only minutes to as much as 2 hours.

Characteristics of attempts in the second stage of labour:

- strength of contractions - moderate (strong),
- duration of contractions - 55-60 seconds,
- frequency - in 1-1,5 minutes.

When the fetus passing through the labour canal its presenting part performs not only translational, but rotational motions. A presenting part of the fetus stretches the pudendal slit, perineum protrudes. In the moment of the highest strain of prelum muscles contractions insertion point of the fetal head – small fontanel is seen from the pudendal slit; between labour pains the head hides in the pudendal slit, and during the next prelum muscles contractions it appears again (*insertion of the head*).

Insertion of the fetal head begins after finishing its internal turn. During each prelum muscles contractions the greater part of the head inserts, and in some time it does not disappear in the pudendal slit in the pause between the prelum muscles contractions.

Disengagement of the fetal head corresponds to the fifth moment of labour's biomechanism — extension of the head. Occipital part of the head disengages for the first time, then — parietal tubers do. Tension of the perineum in the moment of disengagement of the parietal tubers is maximal.

After disengagement of the occiput and parietal tubers forehead and face of the fetus disengage, i. e. the whole head is delivered; during the anterior kind it is turned by its face to the back. During the next strain the internal rotation of the trunk, which corresponds with the external rotation of the head by the face, turned to the mother's hip (in the first position - to the right, in the second - to the left), occurs.

7.3. The third stage of labour comprises that time period between delivery of the baby and delivery of the placenta and may take up to 30 minutes, although usually is much shorter.

The placenta can discharge by fetal (by Schultz) or maternal (by Duncan) surface.

Active management of the postnatal period is used when bleeding occurred and the blood loss is 250 – 300 ml (0,5 % of body weight) while the signs of placental detachment are absent. Active means (manual separation and expulsion of the

afterbirth) are used during inconsiderable external blood loss and in connection with the aggravation of the woman's condition. Attempts to intensify the process of expulsion of the afterbirth by the massage of the uterus, pulling the umbilical cord are excluded, because they break the physiological process of placental detachment from the uterine wall, change the rhythm of its contractions and intensify the bleeding.

8. Management of Normal Labour:

- initial patient evaluation;
- evaluation the risk of maternal and perinatal pathology;
- determine the plan of delivery;
- monitoring of mother and fetus during labour with driving partogram;
- free position of women of the first stage of labour;
- obstetric analgesia and anesthesia;
- newborn care.

8.1. Management of the first stage of labour

8.1.1. External methods for assessing the degree of opening of the cervix.

The speed of moving of the fetal head increases as the result of extension of the dilation of cervix and is usually 1 cm/h in nulliparous women and 2 cm/h - in multiparous, and after the complete cervical dilation - 4 cm/h at the I stage of labour. During the physiological course of labour till the moment of complete cervical dilation inferior pole of the fetal head locates, as a rule, in the narrow part of the pelvic cavity.

According to the data of foreign literature, the position of the fetal head is determined by its correlation to the interspinal line (level of the ischiac spines) - position "0"

8.1.2. Internal methods of assessing the degree of opening of the cervix.

To determine the dynamics of the opening of the cervix and the location of the head of the fetus during labour performed internal obstetrical examination every 4 hours during the first stage of labour and after rupture of membranes.

Indications internal obstetrical examination:

1. Every 4 hours during the first stage of labour and every hour in the second stage of labour;
2. After ruptured of membranes;
3. Pathologic rate of the fetal heart (less than 100 and more than 180 beats / min) to determine the causes that might explain the signs of his condition;
4. Loss umbilical cord;
5. In multiple pregnancies after birth first fetus;

6. Malposition of the fetus;
7. Operative vaginal delivery;
8. Bleeding after 22 weeks gestation (in the operating conditions).


8.1.3. External obstetrical examination

Maneuvers of external obstetrical examination (Leopold's maneuvers).

Evaluation of the extent of the head fitting into the pelvic inlet during labour by palpation is performed with the fourth Leopold's method. If the fingers of the hands can be placed under the head of the fetus and their end touch one another, it locates in the pelvic inlet, if they don't — the head is pressed to its inlet. If the vertex part of the head overhangs the pelvic inlet by 2 cm, and facial part is palpated — the head is put to the pelvis by the minor segment. If the occipital part of the head can not be palpated in the pelvic inlet, and the facial one protrudes by 2–3 cm above the pelvic inlet, the head is put into the major segment. If the chin is palpated or it is not detected, the head is in the pelvic cavity.

Head above brim (as per abdominal palpation):

- 5/5 - The head is mobile over the input to the pelvis;
- 4/5 - The head is pressed against the input to the pelvis;
- 3/5 - The minor segment of the head is in the input to the pelvis;
- 2/5 - The major segment of the head is in the input to the pelvis;
- 1/5 - 0/5 - The head is in the pelvic cavity.



Picture 1. Stations of the fetal head

8.1.4. Determination position of the fetal by internal obstetrical examination

The head is mobile over the input to the pelvis – if the doctor's fingers can be brought under the head.

The head is pressed against the input to the pelvis – if the ends of the doctor's fingers do not meet under the head, nevertheless the occiput and facial parts are palpated above the input to the pelvis.

The minor segment of the head is in the input to the pelvis – the occipital part of the head is over the input to the pelvis by two fingers, and the facial part - completely.

The major segment of the head is in the input to the pelvis – the occipital part of the head is not palpated over the input to the pelvis, and the facial part is over it by two - three fingers

The head is in the pelvic cavity – when only the chin is palpated or no parts of the fetal head are palpated.

Methods of external examination also include measurement of the external sizes of the pelvis. It is performed during the patient's first visit (examination) at the female consultation and at the maternity hospital. If necessary, measurement of the pelvis is repeated during labour.

8.1.5. Cervical Examination

The level - or station - of the presenting fetal part in the birth canal is described in relationship to the ischial spines, which are halfway between the pelvic inlet and the pelvic outlet. When the lowermost portion of the presenting fetal part is at the level of the spines, it is designated as being at zero (0) station. In the past, the long axis of the birth canal above and below the ischial spines was arbitrarily divided into thirds by some and into fifths (approximately 1 cm) by other groups. Adopted the classification of station that divides the pelvis above and below the spines into fifths. Each fifth represents a centimeter above or below the spines. Thus, as the presenting fetal part descends from the inlet toward the ischial spines, the designation is -5, -4, -3, -2, -1, then 0 station. Below the spines, as the presenting fetal part descends, it passes +1, +2, +3, +4, and +5 stations to delivery. Station +5 cm corresponds to the fetal head being visible at the introitus.

If the leading part of the fetal head is at 0 station or below, most often the fetal head has engaged - thus, the biparietal plane has passed through the pelvic inlet. If the head is unusually molded or if there is an extensive caput formation or both, engagement might not have taken place although the head appears to be at 0 station (linea interspinalis).

- mark "-" head is above the linea interspinalis (near the pelvic plane of inlet).
- mark "+" the fetal head is below the linea interspinalis (near the pelvic outlet).

Position of the fetal head in the pelvic plane:

- 3 - head of the fetus above the inlet;
- 2 - head of the fetus pressed to the inlet;
- 1 - head of the fetus by the minor segment in the inlet;
- 0 - head of the fetus by the major segment in the inlet;
- +1 - head of the fetus in the wide part of the small pelvis;
- +2 - head of the fetus in the narrow part of the small pelvis;
- +3 - head of the fetus in the pelvic outlet

8.1.6. Methods of examining the functional condition of the fetus.

Ultrasound fetometry of the fetus is informative after 20 weeks of pregnancy (A) and includes determining the size of the head, circumference of the abdomen and length of the hip. While determining, if there is a discrepancy of one or several basic photometric indicators for the term of pregnancy, an expanded fetometry is performed, where the relation between the biparietal diameter and frontooccipital diameters of the fetal head, size of the head, circumference of the abdomen, biparietal length of the hip bone, hip

bone and circumference of the abdomen are determined. The most valuable indicator is the estimated fetal weight. The basis for performing an ultrasound fetometry in late terms of the pregnancy is suspicion of fetal growth retardation according to external measurements of the height of the uterine fundus and circumference of the woman's abdomen, and also, in special cases, to determine the term of the pregnancy with special tables, if the term is difficult to determine from anamnesis data.

- **Auscultation of the fetal heart tones** is performed after 20 weeks of pregnancy with the help of obstetrical stethoscope, where the frequency of heart beats in one minute is determined.

- physiological normal - 110-170 bpm
- frequency of heart beats above 180 bpm and less than 100 bpm testifies of disorders in the fetal condition.

For auscultation of fetal heart beat use the following rules:

- for facial presentation – listen for the heart beat below the navel on the side where the fetal thorax is located (if first position - on the right side, if second - on the left side).
- for transverse lie - near the navel, closer to the fetal head.
- for breech presentation - above the navel, near the fetal head on the side where the back is turned.

Cardiotocography (CTG) - synchronous electronic monitoring of the fetal heart rate and uterine contractions for 10-15 minutes.

- during analysis of the CTG, such parameters are evaluated: basal frequency of heart rate (BFHR), variability of the heart rate (amplitude and frequency oscillation), presence and type of changes in BFHR in the form of acceleration or decelerations of heart rate.

- if any pathological parameters of heart rate are present, which testify of a dangerous fetal condition, continuous monitoring with the CTG during labour is recommended

- diagnostic criteria: during normal fetal condition for CTG it is characteristic: BFHR is between 110-170 b.p.m (normocardia), variability (width of the tape) - 10-25 bpm with frequency of oscillation 3-6 cycles per minute (wavy type), presence of accelerations and absence of decelerations.

Biophysical profile of the fetus (BPF) (at 30 weeks of pregnancy) – complex evaluation of the condition of the fetus and the sum of separate biophysical parameters (fetal respiratory movements, fetal tone, fetal activity, reactance of the fetal heart rate to non-stress test (NST), volume of amniotic fluid)

- **Possible rupture of membranes.** In 10% of pregnancies, rupture of the membranes precedes the onset of labour. This presents as fluid leaking through the cervix and out of the vagina. The differential diagnosis includes urine leakage, vaginal infections, and

passage of cervical mucus. Because prolonged rupture of the membranes is associated with higher rates of maternal and neonatal infection, optimal treatment of ruptured membranes at term is prompt induction of labour.

8.1.7. Monitoring the condition of women:

- heart rate and blood pressure (every 2 hours)
- temperature (every 4 hours)
- urine: volume; presence of protein or acetone - for displays (every 4 hours).

8.2. Management of the second stage of labour:

- measurement blood pressure, heart rate in women during labour every 10 minutes;
- monitoring of fetal palpitation every 5 minutes during the early phase;
- control by promoting fetal head through the birth canal;
- perform amniotomy if there is no timely rupture of membrane.

Physiologic position and movement.

8.3. Management of the third stage of labour

Two tactics for conducting the third period of delivery exist: *active and conservative*.

Immediately following delivery of the baby, the uterus begins the process of involution. Uterine contractions cause shearing of the placenta away from the uterine wall, and the placenta generally delivers shortly after the baby. Signs of spontaneous placental separation include an apparent lengthening of the umbilical cord, a gush of vaginal bleeding, and a change in shape of the uterus from discoid to globular.

Active management of the third stage of labour has been shown to be of benefit in reducing postpartum blood loss and may include draining the placenta of blood, controlled cord traction, or administration of oxytocic agents. If cord traction is employed, suprapubic pressure with the abdominal hand will lessen the potential for uterine inversion and catastrophic hemorrhage and shock. If the placenta has not delivered within 30 minutes of childbirth, or in the case of severe hemorrhage, the placenta should be manually removed.

The placenta should always be carefully inspected for abnormalities of cord insertion, confirmation of a three-vessel cord, and completeness of removal of the placenta and membranes. If any portion of the placenta or the membranes is missing, the uterine cavity should be manually explored. The uterus should be frequently palpated following delivery of the placenta to ensure that it remains well contracted. Oxytocin, administered as a dilute intravenous solution or given 10 to 20 U intramuscularly, decreases the incidence of postpartum hemorrhage due to uterine atony. The birth canal, including the cervix, vagina, and perineum, should be inspected for lacerations requiring repair. Under most circumstances, the baby can remain with

the mother or immediate family and attempts at breast-feeding within the first 10 to 20 minutes should be encouraged. This first suckling stimulates endogenous oxytocin release and begins the process of milk production and successful breast-feeding.

Episiotomy is an incision in the perineum made to facilitate vaginal delivery. There is no role for routine episiotomy in modern obstetric practice, although there are some clinical indications for its use. In general, episiotomy is used to shorten the second stage of labour for fetal indications (terminal bradycardia or shoulder dystocia) or to control perineal damage when the risk of significant spontaneous laceration is high (operative vaginal delivery, previous large laceration, small perineal body, or large infant). Episiotomy should be performed with adequate local or regional anesthesia and with the verbal consent of the patient, when possible. There are two types of episiotomy techniques in common use: median and mediolateral.

8.3.1. Active conduction of the third stage

Because of a number of advantages, active conduction of the third stage of labour is the most widespread tactic and approved by the World Health Organization, International Federation of Obstetricians-Gynecologists and the International Confederation of Obstetricians.

The use of active conduction of the third stage during each labour lowers the frequency of postnatal bleeding 60% of the time caused by atonia of the uterus, and it also reduces the amount of postnatal blood loss and need for haemotransfusion.

Standard components for active conduction of the third stage include:

- introduction of uterotonics:
- birth of the placenta by controlled traction of the umbilical cord while holding the fundus of the uterus with the palm of the doctor's hand;
- massage of the uterus through anterior abdominal wall after the birth of the placenta.

Rules for introducing uterotonics: within the first minute after the birth of the child palpate the uterus for the presence of a second child, if there is no other child – introduce 10 units of oxytocin i/m. Oxytocin is the most widespread uterotonic because it takes effect in 2-3 minutes; it can be used for all women.

If oxytocin can not be used, use ergometrin - 0,2 mg i/m. The woman should be informed about the possible side-effects of these preparations.

Ergometrin cannot be used in women with pre-eclampsia, eclampsia and hypertension.

Controlled traction by the umbilical cord:

- clamp the umbilical cord closer to the perineum; hold the clamped umbilical cord and clamp in one hand;

- put the second hand directly over the women's pubis and hold the uterus, pulling away from the symphysis;
- slightly pull the umbilical cord and wait for a strong contraction of the uterus (usually 2-3 minutes after the introduction of oxytocin);
- simultaneously during the strong contraction, the woman should push and very cautiously pull (traction) the umbilical cord downward till the birth of the placenta; simultaneously continue with the second hand contraction in the opposite direction of traction (pushing the uterus away from the symphysis).
- if the placenta does not detach during 30-40 seconds of controlled traction, stop the traction by the umbilical cord, but continue cautiously keeping the cord in light tension; the second hand remains over the pubis, holding the uterus.
- wait for the uterus to contract again and repeat the controlled traction by the umbilical cord with contraction of the uterus.

Never use traction (pulling downwards) by the umbilical cord without contraction of a well contracted uterus over the pubis.

Using traction by the umbilical cord without contraction of the uterus can lead to prolapse or inversion of the uterus.

After the placenta is delivered, hold it with both hands and cautiously turn it, pulling the membranes out. If the membranes tear, cautiously examine the vagina and cervix in sterile gloves. If the membranes are seen, carefully use a clamp to remove it.

Attentively examine the placenta and make certain of its integrity. If an area of the maternal surface is absent, or if there is an area torn with vessels, there is reason to suspect retention of an area of the placenta and begin necessary measures.

Massage of the uterus: after the birth of the placenta immediately massage the uterus through the anterior abdominal wall until the uterus does not become firm.

Further, the uterus should be palpated every 15 minutes for the first 2 hours, to be sure that after the uterus is massaged it does not relax, but remains firm. If necessary repeat the massage.

Ice is not applied on the lower abdomen during the early postnatal period.

Active conduction of the III stage of labour should be offered to each woman as it lowers the frequency of postnatal bleedings resulting from atonia of the uterus. The parturient woman should be informed concerning active conduction of the III stage of labour, and should give voluntary written consent.

8.3.2. Passive conduction of the third stage of labour

The postnatal period of labour is the shortest (5 - 30 min). However, very important because of the possibility of appearing of postnatal bleeding. The postnatal period is accompanied by physiological blood loss (0,5 % of the woman's weight).

The midwife, when the umbilical cord stops pulsating, but no later than one minute after the birth of the child, clamps and cuts the umbilical cord. The general condition of the woman is carefully supervised; signs of placental detachment, amount of blood loss are closely watched.

When signs of placental detachment occur (Schreder's sign, Alfred's, Klein's, Kustner-Chukalov's) - it is necessary to have the woman "push" which leads to the birth of the afterbirth.

If there are no signs of placental detachment or signs of external bleeding 30 minutes after the delivery of the baby, manual detachment and delivery of the afterbirth is performed. If there are signs of bleeding - manual detachment and delivery of the afterbirth should be performed immediately with adequate anesthesia.

After the placenta is delivered, it should be carefully examined (be certain of the integrity of the placenta and membranes).

The general duration of birth on average for primipara is 8-12 hours, for secundipara - 6-8 hours.

The birth canal is examined after the delivery (with the help of vaginal mirrors) only if there is excessive bleeding, after operative vaginal delivery or if the doctor is uncertain about the integrity of the birth canal (fast childbirth, childbirth outside the hospital).

9. Partogram

The partogram provides a graphical illustration of the progress of labour and is considered by the World Health Organisation (WHO) to be a valuable tool for managing intrapartum women.

The partogram:

- Depicts the progress of labour at a glance,
- Enables failure to progress to be readily recognised,
- Is simple to use,
- Provides a practical teaching aid,
- Is an efficient means of exchange of technical information about labour progress between teams of caregivers.

KEY POINTS:

1. The partogram should be used for all women admitted in established labour. When the partogram is commenced at the beginning of the induction process the Alert and Action lines are drawn when the woman is in the active phase of labour.

2. Established labour is defined as the presence of regular contractions, increasing in strength and duration, leading to progressive effacement and dilatation of the cervix.
3. A rate of 1cm/hour in the active phase of labour is often accepted as normal progress in labour. Many women who show slower rates of cervical dilation will proceed to normal birth.²
4. The Alert line is a simple tool which separates women into two groups:
Women with cervical dilatation equal to / greater than 1cm/hour who are highly unlikely to require operative intervention.
Women with cervical dilatation slower than 1 cm/hour who are more likely to require operative intervention.
5. The WHO partogram does not differentiate between nulliparous or multiparous women's labours.

MANAGEMENT:

A vaginal examination is performed 4 hours after the initial one or earlier if clinically warranted. If subsequent examination shows dilatation between Alert line and Action line a repeat vaginal examination is carried out in 2 hours. At this examination if the cervical dilatation is touching / crossing the Action line, the Labour and Birth Suite medical team must evaluate the woman's progress in labour and instigate appropriate intervention.

ANNEX 2: Partograph




Fig. 4. Recording progress of labour.

10. Newborn Care. Immediate Assessment

In 1952, Dr. Virginia Apgar devised a 0 to 10 scale scoring system intended to focus attention on the newborn and allow the systematic assessment of its condition and immediate needs. Since that time, Apgar scores have become a mainstay in immediate newborn assessment.

To prevent the aspiration mucus is drained from the nasal part of the pharynx of the newborn immediately after the external rotation of the head, when its chest is compressed by the maternal passages and he can not make the first inhalation. After the first inhalation skin of the newborn becomes rose, he begins to move by his extremities, loudly screams.

Condition of the newborn is estimated by Virginia Apgar's score on the 1st and 5th min of his life.

Your baby will be checked for five things:

- A – Activity; muscle tone
- P – Pulse rate
- G – Grimace; reflex (measured by placing a bulb syringe in the baby's nose and seeing his response)
- A – Appearance; skin color
- R – Respiration

Each category is given a score ranging from 0-2. The numbers are then added up for a final score.

By the Apgar's score the majority of newborns on the 1st minute of life has 7 - 8 points (acrocyanosis is detected as the result of transitional blood circulation and decrease in the muscular tonus); 4 - 6 points asphyxia of slight degree; 0 - 3 - severe asphyxia.

The mark 10 points on the 1st minute of life is detected only in 15% of the newborns. On the 5th minute of life estimation by the Apgar's score in the norm rises up to 8 - 10 points.

11. Newborn Care

After 1 minute babies born:

- After the stop of pulsation of the umbilical cord two clamps are applied on it (10 cm from the umbilical ring), processed by 5% alcoholic solution of iodine and incised by aseptic scissors. During the physiological labour it is recommended to put a newborn to the breast of the mother (within first 2 h), better till the section of the umbilical cord, which contributes to its better adaptation and reduce of the length of postnatal stage of labour.

- Keep baby warm. When a baby is born, he is wet from the fluid in the womb and can easily become cold. Nurses will dry his skin, wrap him in a blanket, place a knitted hat on his head and may even use heat lamps to help him stay warm. Holding your baby close to you so that your skin touches his also helps keep him warm.

After the labour of the newborn its height (from pate to hill), weight, circumference of the head (direct diameter) and shoulders are measured. Bracelets are put on, where the sex, mother's surname and name, number of labour history, weight and height, date and time of delivery are written. The newborn is covered with warm wrapping clothes and blanket and put on the warm wrapping table.

12. Anaesthesia and analgesia of labour.

- Physiologic positions and movements,
- Child birth preparation: Psychoprophylaxis,
- Hydrotherapy• Hydrotherapy involve a simple shower or tubbath, or it include the use of a whirlpool or large tub specially equipped for pregnant patients. Benefits of

hydrotherapy includes reduced pain & anxiety, decreased BP & increased efficiency of uterine relaxation.

- Systemic Medication for Labour Analgesia

Obstetric anesthesia encompasses all techniques used by anesthesiologists and obstetricians to alleviate the pain associated with labour and delivery: general anesthesia, regional anesthesia, local anesthesia, and analgesia. Relief from pain during labour and delivery is an essential part of good obstetric care. Unique clinical considerations guide anesthesia provision for obstetric patients; physiologic changes of pregnancy and increases in certain complications must be considered.

Obstetric analgesia or anesthesia refers to the multiple techniques useful for the alleviation of pain associated with labour, delivery, or surgery. The choice of an appropriate analgesic technique must be made by the patient, the obstetrician, and the anesthesiologist and should take into consideration the patient's anatomy and physiology, the status of her fetus, the obstetric plan for delivery, and the pharmacology of the drugs to be employed.



The Notorious K.I.M.'s Physiologic Positions & Movements For Labor And Birth

V. Supportive employment.

3.3. Requirements for the results of work.

1. To know the precursors birth, preliminary period.
2. To give the evaluation of the cervical status (scale Bishop).
3. To show the biomechanism of labour at the anterior and posterior types occipital presentation.
4. To tell the clinical course of the first stage of labour. Methods registration labour. Monitoring during delivery.
5. To name periods of delivery and tell clinical management of the first stage of normal labour. The basic principles of partogram.
6. To perform the clinical course of the second stage of labour and clinical management of this stage of labour. To perform the third stage of labour. The signs of placental separation. Clinical management III stage of labour (active and expectant tactics).
7. Delivery of the placenta according to Abuladze, Genter's method and Crede-Lazarevich's method.
8. To show the examination of the placenta. Physiological and pathological hemorrhage.
9. To assess the essentials of normal newborn assessment (Apgar scores) and care. Care and observation in the first few hours of life.
10. To tell about obstetric analgesia and anesthesia.

ALGORITHM for conducting normal labour:

- evaluate the degree of predicted risk of development of maternal and perinatal pathologies so as to determine the necessary level of help during labour;
- determine the plan for conducting labour and an informed conformation from the woman;
 - emotional support for the parturient woman during labour (labour with partner);
 - control over the condition of the mother and fetus during labour while conducting the partogram;
 - free position of the parturient woman during labour;
 - pain relief if indicated;
 - condition of the fetus:
- frequency of fetal heart beats, performed with auscultation or manual Doppler analyzer (every 15 minutes)
- degree of configuration of the fetal head (every 4 hours)
- condition of the amniotic sac and amniotic fluid (every 4 hours)
 - condition of the parturient woman:

- pulse and arterial pressure (every 2 hours)
- body temperature (every 4 hours)
- urine: amount; presence of protein or acetone – if indicated (every 4 hours).
 - conducting the second period of labour:
- measurement of the woman's BP, pulse every 10 minutes;
- control of fetal heart activity every 5 minutes during the early phase, and after each contraction during the active phase;
- control of the advancement of the fetal head through the birth canal by using internal obstetrical examination each hour.

Because of increased risk of ascending infection through the birth canal additional internal obstetrical examinations during the second stage of labour are admissible only if indicated:

- performing amniotomy if the amniotic sac has not ruptured
- during multiple pregnancy after the birth of the first baby
- when deciding on operative vaginal delivery (obstetrical forceps, vacuum extraction, extraction of the fetus by the pelvic end).
- evaluate the condition of the child during birth; perform primary care of the newborn and early breast feeding, principles of "thermal chain"
 - Preventive measures are taken for ophthalmi for all newborns during the first hour of life; 0,5% erythromycin or 1 % tetracycline ointments are used.
 - 2 hours after the birth, the baby is changed into clean clothes and together with the mother is taken to the postnatal ward.

3.4. Control materials for the final stage of the class: tasks, tests, etc.

Tests

1. The head of a newborn has a dolichocephalic form, extended from the front to the back. At the occipital area, there is a patrimonial tumour located in the middle between the large and small fontanel. What head presentation did the birth occur in?
 - A. Frontoparietal.
 - B. Anterior type, occipital presentation.
 - C. Posterior type, occipital presentation.
 - D. Facial.
 - E. Frontal.
2. Primipara, 20 years old, is in the beginning of the I period of physiological birth. Contractions are 15-20 sec., every 10-15 min., weak. Fetal palpitation is not suffering. How much in cm should the cervix be open for the amniotic discharge to be timely?
 - A. 6-8.
 - B. 8-10.
 - C. 4-6.

- D. 2-4.
E. 1,5-2.
3. During internal obstetrical examination of a secundipara, the cervix is open 4-5 cm, when pressing with a finger on the fetal head, it pushes away and again returns to its initial position, the pelvis is free. How long is the fetal head permitted to be in this plane, in hours?
A. 5.
B. 6.
C. 4.
D. 3.
E. 2.
4. You are observing a parturient woman in the 1 period of birth. During internal obstetrical examination, the sagittal suture is in the right diagonal diameter, the small fontanel is closer to the pubis. How should the parturient woman lay in bed, so that the fetal head would be insert correctly?
A. On her left side.
B. On her back.
C. On her right side.
D. It make do difference.
E. She is only allowed to walk.
5. Secundipara with a weight of 80 kg. What is the acceptable amount of blood loss, ml:
A. 400.
B. 500.
C. 600.
D. 700.
E. 800.
6. A woman in her first pregnancy had a live baby boy, weight 3200 g, length 50 cm. The umbilical cord was transected after pulsation of the vessels stopped. When the edge of the palm is pressed on the symphysis, the umbilical cord retreats into the vagina. What sign is used for determining whether the placenta has separated from the uterus?
A. Alfelda.
B. Kustner-Chukalova.
C. Schröder.
D. Dovjenko
E. Rogovina.

Answer key

1.	B	7.	
2.	B	8.	
3.	B	9.	
4.	C	10.	
5.	A	11.	
6.	B	12.	

4. SUMMING UP

Assessment of the ongoing learning activity at the practical class:

1. Assessment of the theoretical knowledge on the theme:
 - methods: individual survey on the theme, participation of the students in the discussion of problem situations; assessment of performance of tests on the theme;
 - the maximum score – 5, the minimum score – 3, the unsatisfactory score – 2.
2. Assessment of practical skills on the theme:
 - methods: assessment of the solution of situational tasks (including calculation) on the theme;
 - the maximum score – 5, the minimum score – 3, the unsatisfactory score – 2.

Assessment of the individual task:

1. Assessment of the quality of the performance of the individual task:
 - the maximum score – 5, the minimum score – 3, the unsatisfactory score – 2.
2. Assessment of the presentation and defense of an individual task, participation in the assessment of the business plan of the competitors and its critical analysis:
 - the maximum score – 5, the minimum score – 3, the unsatisfactory score – 2.

The score for one practical class is the arithmetic average of all components and can only have an integer value (5, 4, 3, 2), which is rounded statistically.

Criteria for ongoing assessment at the practical class:

5	The student is fluent in the material, takes an active part in the discussion and solution of situational clinical problems, confidently demonstrates practical skills during the examination of a pregnant women and woman who is giving birth and interpretation of clinical, laboratory and instrumental studies, expresses his opinion on the topic, demonstrates clinical thinking.
4	The student is well versed in the material, participates in the discussion and solution of situational clinical problems, demonstrates practical skills during the examination of a pregnant and just delivered woman and

	interpretation of clinical, laboratory and instrumental studies with some errors, expresses his opinion on the topic, demonstrates clinical thinking.
3	The student isn't well versed in material, insecurely participates in the discussion and solution of a situational clinical problem, demonstrates practical skills during the examination of a pregnant or woman during delivery or any period of labour and interpretation of clinical, laboratory and instrumental studies with significant errors.
2	The student isn't versed in material at all, does not participate in the discussion and solution of the situational clinical problem, does not demonstrate practical skills during the examination of a pregnant or woman in any period of labour or in early after delivery period and the interpretation of clinical, laboratory and instrumental studies.

RECOMMENDED LITERATURE

Basic:

1. Gladchuk I.Z. Obstetrics: student`s book / Gladchuk I.Z., Ancheva I.A. . – Vinnitsia: Nova Knyha, 2021. – 288 p.
2. Obstetrics and Gynecology: in 2 volumes. Volume 1. Obstetrics: textbook / V.I. Gryshchenko, M.O. Shcherbina, B.M. Ventskiivskyi et al. (2nd edition). – «Medicina», 2018. – 392 p.
3. Hiralal Konar DC Dutta's Textbook of Obstetrics (9th Ed.) / Hiralal Konar (Ed.). – Jp Medical Ltd, 2018. – 700 p.
4. F. Gary Cunningham Williams Obstetrics (26th Edition) / F. Gary Cunningham, Kenneth Leveno, Jodi Dashe, Barbara Hoffman, Catherine Spong, Brian Casey. – McGraw Hill / Medical, 2022. – 1328 p.
5. Jeremy Oats, Suzanne Abraham Llewellyn-Jones Fundamentals of Obstetrics and Gynaecology (10th Ed) / Jeremy Oats, Suzanne Abraham. – Elsevier, 2016. – 384 p.

Additional:

1. The PROMPT-CIPP Editorial Team. (2019). PROMPT-CIPP Course Participant's Handbook: Care of the Critically Ill Pregnant or Postpartum Woman (Critical Car Prompt Practical Obstetric Multi-professional Training). – Cambridge University Press; 1st edition, 2019. – 136 p.
2. L. A. Magee The FIGO Textbook of Pregnancy Hypertension. An evidence-based guide to monitoring, prevention and management. / L. A. Magee, P. Dadelszen, W. Stones, M. Mathai (Eds). – The Global Library of Women's Medicine, 2016. – 456 p.

3. Edwin Chandrachan Handbook of CTG Interpretation: From Patterns to Physiology / Edwin Chandrachan. – Cambridge University Press; 1st edition, 2017. – 256 p.
4. Louise C. Kenny, Jenny E. Myers Obstetrics by Ten Teachers (20th ed) / Louise C. Kenny, Jenny E. Myers. – CRC Press, 2017. – 342 p.
5. J. Studd Current Progress in Obstetrics and Gynaecology. Vol 4. / J. Studd, Seang Lin Tan, F. Chervenak. – TreeLife Media (A Div of Kothari Medical), 2017. – 419 p.
6. J. Studd Current Progress in Obstetrics and Gynaecology. Vol 5. / J. Studd, Seang Lin Tan, F. Chervenak. – TreeLife Media (A Div of Kothari Medical), 2019. – 403 p.
7. J. Studd Current Progress in Obstetrics and Gynaecology. Vol 6. / J. Studd, Seang Lin Tan, F. Chervenak. – TreeLife Media (A Div of Kothari Medical), 2022. – 309 p.
8. Mark Landon Obstetrics: Normal and Problem Pregnancies, 8th Edition / Mark Landon, Henry Galan, Eric Jauniaux, Deborah Driscoll, Vincenzo Berghella, William Grobman, et al. – Elsevier, 2021. – 1280 pp.
9. Mark B. Landon Gabbe's Obstetrics Essentials: Normal & Problem Pregnancies, 1st Edition / Mark B. Landon, Deborah A. Driscoll, Eric R. M. Jauniaux, Henry L. Galan, William A. Grobman, Vincenzo Berghella. – Elsevier, 2019. – 496 pp.
10. Ian M. Symonds, Sabaratnam Arulkumaran Essential Obstetrics and Gynaecology, 6th Edition / Ian M. Symonds, Sabaratnam Arulkumaran. – Elsevier, 2020. – 480 pp.
11. Myra J. Wick Mayo Clinic Guide to a Healthy Pregnancy, 2nd Edition / Myra J. Wick. – Mayo Clinic Press, 2018. – 520 p.

INTERNET SOURCES:

- <https://www.cochrane.org/>
- <https://www.ebcog.org/>
- <https://www.acog.org/>
- <https://www.uptodate.com>
- <https://online.lexi.com/>
- <https://www.ncbi.nlm.nih.gov/>
- <https://pubmed.ncbi.nlm.nih.gov/>
- <https://www.thelancet.com/>
- <https://www.rcog.org.uk/>
- <https://www.npwh.org/>